# **Complete Summary**

#### **TITLE**

Diagnosis and treatment of respiratory illness in children and adults: percentage of patients with a diagnosis of pharyngitis who had strep screen testing.

## SOURCE(S)

Institute for Clinical Systems Improvement (ICSI). Diagnosis and treatment of respiratory illness in children and adults. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2008 Jan. 71 p. [175 references]

### **Measure Domain**

#### **PRIMARY MEASURE DOMAIN**

**Process** 

The validity of measures depends on how they are built. By examining the key building blocks of a measure, you can assess its validity for your purpose. For more information, visit the <u>Measure Validity</u> page.

#### **SECONDARY MEASURE DOMAIN**

Does not apply to this measure

## **Brief Abstract**

### **DESCRIPTION**

This measure is used to assess the percentage of patients with a diagnosis of pharyngitis who had strep screen testing.

## **RATIONALE**

The priority aim addressed by this measure is to reduce excessive antibiotic treatment through decreased empiric treatment of patients with pharyngitis.

#### PRIMARY CLINICAL COMPONENT

Pharyngitis; rapid strep test; strep culture

#### **DENOMINATOR DESCRIPTION**

Number of patients with a diagnosis of pharyngitis as identified by the following International Classification of Diseases, Ninth Revision (ICD-9) codes: 462.0 and 034.4

#### **NUMERATOR DESCRIPTION**

Number of patients with a diagnosis of pharyngitis who received a strep screen test\*

## **Evidence Supporting the Measure**

## **EVIDENCE SUPPORTING THE CRITERION OF QUALITY**

 A clinical practice guideline or other peer-reviewed synthesis of the clinical evidence

#### NATIONAL GUIDELINE CLEARINGHOUSE LINK

• <u>Diagnosis and treatment of respiratory illness in children and adults.</u>

## **Evidence Supporting Need for the Measure**

## **NEED FOR THE MEASURE**

Unspecified

## **State of Use of the Measure**

#### **STATE OF USE**

Current routine use

## **CURRENT USE**

Internal quality improvement

# **Application of Measure in its Current Use**

### **CARE SETTING**

Physician Group Practices/Clinics

## PROFESSIONALS RESPONSIBLE FOR HEALTH CARE

**Physicians** 

<sup>\*</sup>Rapid strep test or strep culture

## LOWEST LEVEL OF HEALTH CARE DELIVERY ADDRESSED

**Group Clinical Practices** 

#### **TARGET POPULATION AGE**

Age greater than or equal to 3 months

#### **TARGET POPULATION GENDER**

Either male or female

#### STRATIFICATION BY VULNERABLE POPULATIONS

Unspecified

## **Characteristics of the Primary Clinical Component**

## INCIDENCE/PREVALENCE

After viral upper respiratory infection and otitis, acute pharyngitis and/or tonsillitis is the third most common illness diagnosed by U.S. pediatricians.

Group A beta streptococcal pharyngitis is uncommon in children younger than three years of age and rare in children younger than 18 months old.

## **EVIDENCE FOR INCIDENCE/PREVALENCE**

Institute for Clinical Systems Improvement (ICSI). Diagnosis and treatment of respiratory illness in children and adults. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2008 Jan. 71 p. [175 references]

## **ASSOCIATION WITH VULNERABLE POPULATIONS**

Unspecified

## **BURDEN OF ILLNESS**

Unspecified

## **UTILIZATION**

Unspecified

## **COSTS**

Unspecified

## **Institute of Medicine National Healthcare Quality Report Categories**

#### **IOM CARE NEED**

**Getting Better** 

#### **IOM DOMAIN**

Effectiveness

## **Data Collection for the Measure**

#### **CASE FINDING**

Users of care only

## **DESCRIPTION OF CASE FINDING**

Patients with pharyngitis diagnosis.

Identify patients with pharyngitis diagnosis of 462.0 and 034.4. The medical record of each patient is reviewed to determine if the patient had either a rapid strep test or strep culture.

The suggested time period for data collection is a calendar month.

## **DENOMINATOR SAMPLING FRAME**

Patients associated with provider

## **DENOMINATOR INCLUSIONS/EXCLUSIONS**

#### **Inclusions**

Number of patients with a diagnosis of pharyngitis as identified by the following International Classification of Diseases, Ninth Revision (ICD-9) codes: 462.0 and 034.4

#### **Exclusions**

Unspecified

# **RELATIONSHIP OF DENOMINATOR TO NUMERATOR**

All cases in the denominator are equally eligible to appear in the numerator

# **DENOMINATOR (INDEX) EVENT**

Clinical Condition Encounter

## **DENOMINATOR TIME WINDOW**

Time window is a single point in time

## **NUMERATOR INCLUSIONS/EXCLUSIONS**

#### **Inclusions**

Number of patients with a diagnosis of pharyngitis who received a strep screen test\*

\*Rapid strep test or strep culture

#### **Exclusions**

Unspecified

# MEASURE RESULTS UNDER CONTROL OF HEALTH CARE PROFESSIONALS, ORGANIZATIONS AND/OR POLICYMAKERS

The measure results are somewhat or substantially under the control of the health care professionals, organizations and/or policymakers to whom the measure applies.

## **NUMERATOR TIME WINDOW**

Encounter or point in time

#### **DATA SOURCE**

Administrative data Medical record

### **LEVEL OF DETERMINATION OF QUALITY**

Individual Case

#### **PRE-EXISTING INSTRUMENT USED**

Unspecified

## **Computation of the Measure**

#### **SCORING**

Rate

#### **INTERPRETATION OF SCORE**

Better quality is associated with a higher score

#### **ALLOWANCE FOR PATIENT FACTORS**

Unspecified

#### STANDARD OF COMPARISON

Internal time comparison

# **Evaluation of Measure Properties**

#### **EXTENT OF MEASURE TESTING**

Unspecified

## **Identifying Information**

## **ORIGINAL TITLE**

Percentage of patients with a diagnosis of pharyngitis who had strep screen testing.

#### **MEASURE COLLECTION**

Diagnosis and Treatment of Respiratory Illness in Children and Adults Measures

#### **DEVELOPER**

Institute for Clinical Systems Improvement

# **ADAPTATION**

Measure was not adapted from another source.

## **RELEASE DATE**

2007 Jan

## **REVISION DATE**

2008 Jan

## **MEASURE STATUS**

This is the current release of the measure.

This measure updates a previous version: Institute for Clinical Systems Improvement (ICSI). Diagnosis and treatment of respiratory illness in children and adults. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2007 Jan. 71 p.

## SOURCE(S)

Institute for Clinical Systems Improvement (ICSI). Diagnosis and treatment of respiratory illness in children and adults. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2008 Jan. 71 p. [175 references]

#### **MEASURE AVAILABILITY**

The individual measure, "Percentage of Patients With a Diagnosis of Pharyngitis Who Had Strep Screen Testing," is published in "Health Care Guideline: Diagnosis and Treatment of Respiratory Illness in Children and Adults." This document is available from the <u>Institute for Clinical Systems Improvement (ICSI) Web site</u>.

For more information, contact ICSI at, 8009 34th Avenue South, Suite 1200, Bloomington, MN 55425; phone: 952-814-7060; fax: 952-858-9675; Web site: <a href="https://www.icsi.org">www.icsi.org</a>; e-mail: <a href="https://icsi.info@icsi.org">icsi.info@icsi.org</a>.

## **NQMC STATUS**

This NQMC summary was completed by ECRI Institute on April 4, 2007. The summary was updated by ECRI Institute on June 18, 2008.

#### **COPYRIGHT STATEMENT**

This NQMC summary (abstracted Institute for Clinical Systems Improvement [ICSI] Measure) is based on the original measure, which is subject to the measure developer's copyright restrictions.

The abstracted ICSI Measures contained in this Web site may be downloaded by any individual or organization. If the abstracted ICSI Measures are downloaded by an individual, the individual may not distribute copies to third parties.

If the abstracted ICSI Measures are downloaded by an organization, copies may be distributed to the organization's employees but may not be distributed outside of the organization without the prior written consent of the Institute for Clinical Systems Improvement, Inc.

All other copyright rights in the abstracted ICSI Measures are reserved by the Institute for Clinical Systems Improvement, Inc. The Institute for Clinical Systems Improvement, Inc. assumes no liability for any adaptations or revisions or modifications made to the abstracts of the ICSI Measures.

#### Disclaimer

## **NQMC DISCLAIMER**

The National Quality Measures Clearinghouse™ (NQMC) does not develop, produce, approve, or endorse the measures represented on this site.

All measures summarized by NQMC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public and private organizations, other government agencies, health care organizations or plans, individuals, and similar entities.

Measures represented on the NQMC Web site are submitted by measure developers, and are screened solely to determine that they meet the NQMC Inclusion Criteria which may be found at

http://www.qualitymeasures.ahrq.gov/about/inclusion.aspx.

NQMC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or its reliability and/or validity of the quality measures and related materials represented on this site. The inclusion or hosting of measures in NQMC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding measure content are directed to contact the measure developer.

© 2008 National Quality Measures Clearinghouse

Date Modified: 11/3/2008

